

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

Spatio-temporal Drought Trends by Forest Type in the Conterminous United States, 1960-2013

Matthew P. Peters, Louis R. Iverson, and Stephen N. Matthews


aG3701
.K1
2014
.P48
Map

Map Description

Droughts are common in virtually all U.S. forests, but their frequency and intensity vary within forest ecosystems (Hanson and Weltzin 2000). Accounting for the long-term influence of droughts within a region is difficult due to variations in the spatial extent and intensities over a period. Therefore, we created a cumulative drought severity index (CDSI) (Fig. 1) for the United States using weighted monthly frequencies of Palmer Drought Severity Index values reported at climate divisions for the periods 1960 to 1986 and 1987 to 2013 (National Climate Data Center 2014, Palmer 1965). The occurrences of 'severe' conditions (index value of -3.0 to -3.9) receive a weight of 2 and 'extreme' (index value of ≤ -4.0) receive a weight of 3 to reflect the increased magnitude of these events.

The Authors


Matthew P. Peters is a forestry technician and Louis R. Iverson is a research landscape ecologist with the Northern Research Station in Delaware, OH. Stephen N. Matthews is a research assistant professor in the School of Environment and Natural Resources at Ohio State University and is affiliated with U.S. Forest Service at the Northern Research Station.



Spatio-temporal Drought Trends by Forest Type in the Conterminous United States, 1960-2013

Matthew P. Peters, Louis R. Iverson, and Stephen N. Matthews

1960-1986



1987-2013



Cumulative Drought Severity Index

- < 50
- 51 - 100
- 101 - 150
- 151 - 200
- 201 - 250
- > 250
- U.S. Lakes

Data Sources
National Climatic Data Center 2014
North American Drought Monitor Indices
<http://www.ncdc.noaa.gov/temp-and-precip/drought/nadmi/indices.php?>
U.S. Forest Service and U.S. Geological Survey 2000 Forest Cover Types
<http://www.fia.fs.fed.us/library/maps/>

Map Description

Droughts are common in virtually all U.S. forests, but their frequency and intensity vary within forest ecosystems (Hanson and Weltzin 2000). Accounting for the long-term influence of droughts within a region is difficult due to variations in the spatial extent and intensities over a period. Therefore, we created a cumulative drought severity index (CDSI) (Fig. 1) for the United States using weighted monthly frequencies of Palmer Drought Severity Index values reported at climate divisions for the periods 1960 to 1986 and 1987 to 2013 (National Climatic Data Center 2014, Palmer 1965). The occurrences of 'severe' conditions (index value of -3.0 to -3.9) receive a weight of 2 and 'extreme' (index value of ≤ -4.0) receive a weight of 3 to reflect the increased magnitude of these events.




Figure 1. Cumulative drought severity index values, which are a weighted summation of monthly drought occurrences, are mapped among climate divisions of the conterminous United States for the periods 1960 to 1986 and 1987 to 2013. Forest cover types defined by the U.S. Forest Service were used to mask nonforested land depicting the impact of droughts within forests (large map and inset right).

Forest cover types defined by the U.S. Forest Service include 21 distinct groups that were used to mask nonforested land and summarize weighted frequencies of drought occurring during each 324-month period. Additionally, forest cover types were combined within eastern and western United States and drought frequencies among climate divisions were summarized for the two periods (Figure 2).

Drought severity has increased throughout the West, Southeast, and Lake States (see map at right). The Northeast and Mid-Atlantic States have not experienced such severe droughts in the latter period. The mapped areas show that the frequencies of drought conditions in forests have significantly increased in the West compared to the East.

Drought frequencies for forested land of the conterminous United States during the period 1960-2013 were adjusted for the number months per climate division. For each forest type, drought conditions were summarized as the percentage of months during the 324-month period (27 years) among climate divisions that contained the forested land (Table 1). Data used to create these maps can be obtained by contacting the author: matthewpeters@fs.fed.us.




Figure 2. Comparison of Palmer Drought Severity Index classes among eastern and western U.S. forested land for the period 1960-2013.

Table 1. Percent of drought occurrences among major forest types of the conterminous United States and relative cumulative drought severity index (CDSI) scores. Droughts reported at climate divisions containing each forest type were used to aggregate values and calculate percentages of the 324-month period.

Forest Type	U.S. area (km ²)	1960-1986				1987-2013			
		MD	SD	EO	CDSI	MD	SD	EO	CDSI
Aspen-birch	106,263	8.6	4.0	4.3	29.5	7.5	4.3	2.7	24.2
Elm-ash-cottonwood	21,481	8.5	4.2	3.4	27.1	8.7	5.0	3.5	29.2
Loblolly-shortleaf pine	235,600	10.0	3.9	1.0	20.8	11.5	6.0	2.5	31.0
Longleaf-slash pine	81,888	10.7	3.7	0.7	20.2	11.8	6.1	2.7	32.1
Maple-beech-birch	204,826	8.6	3.9	2.9	25.1	8.7	4.5	2.3	24.6
Oak-gum-cypress	119,125	10.2	4.0	1.0	21.2	11.4	5.9	2.5	30.7
Oak-hickory	501,134	9.3	4.0	2.2	23.9	9.6	5.0	2.5	27.1
Oak-pine	192,825	9.5	3.8	1.4	21.3	10.1	4.9	1.9	25.6
Spruce-fir	58,803	8.7	4.2	3.2	26.7	7.7	3.9	2.2	22.1
White-red-jack pine	71,113	8.9	4.1	3.4	27.3	8.2	4.3	1.9	22.5
Chaparral	52,182	10.8	4.5	2.3	26.7	14.4	12.2	9.7	67.9
Douglas-fir	153,728	9.4	4.5	3.0	27.4	13.2	10.2	9.6	62.4
Fir-spruce	76,688	9.7	4.2	2.5	25.6	13.2	10.3	9.5	62.3
Hemlock-Sitka spruce	17,596	10.9	2.0	1.0	18.2	13.0	5.8	1.6	29.4
Larch	11,439	7.9	4.1	4.0	28.1	15.8	12.5	8.3	65.5
Lodgepole pine	117,398	9.6	4.7	3.7	30.1	12.8	10.2	9.9	62.9
Pinon-juniper	224,678	11.1	4.8	2.0	26.7	13.7	11.0	8.3	60.6
Ponderosa pine	221,476	10.0	4.5	3.1	28.3	15.0	10.2	8.7	59.5
Redwood	5,998	11.1	3.0	1.5	21.6	16.7	6.7	1.3	34.0
Western hardwoods	65,933	9.9	4.3	2.8	26.9	13.6	10.6	8.0	61.5
Western white pine	9,048	7.9	3.5	3.7	28.0	15.1	12.3	10.7	71.8
Eastern Forest	1,593,058	9.3	4.0	2.3	24.2	9.5	5.0	2.5	27.0
Western Forest	956,222	9.8	4.0	2.7	25.9	14.0	10.2	7.9	58.1

MD= moderate drought (-2.0 to -2.9); SD=severe drought (-3.0 to -3.9); EO=extreme drought (≤ -4)

Data Sources

Palmer Drought Severity Index values were obtained from the National Climatic Data Center (2014) and used to calculate the CDSI. The forest cover types layer was provided by the U.S. Forest Service and U.S. Geological Survey (2000) and used to mask nonforested land.

Suggested Citation

Peters, Matthew P.; Iverson, Louis R.; Matthews, Stephen N. 2014. Spatio-temporal trends of drought by forest type in the conterminous United States, 1960-2013. Res. Map NRS-7. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. [scale 1:12,000,000].

Literature Cited

Hanson, P.J.; Weltzin, J.F. 2000. Drought disturbance from climate change: response of United States forests. Science of the Total Environment. 262(3): 205-220.

National Climatic Data Center. 2014. North American drought monitor indices. Asheville, NC: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Climatic Data Center. Available at www.ncdc.noaa.gov/temp-and-precip/drought/nadmi/indices.php? (Accessed July 22, 2014).

Palmer, W.C. 1965. Meteorological drought. Washington, DC: U.S. Department of Commerce. 58 p.

U.S. Forest Service and U.S. Geological Survey. 2000. Forest cover types. Washington, DC: U.S. Department of Agriculture. Available at www.fia.fs.fed.us/library/maps/ (Accessed July 22, 2014).

The USDA is an equal opportunity provider and employer.

Manuscript received for publication
August 2014

Published by
USDA Forest Service
Northern Research Station
11 Campus Blvd., Suite 200
Newtown Square, PA 19073
December 2014

For additional copies:
USDA Forest Service
Publications Distribution
359 Main Road
Delaware, OH 43015
Fax: 740-368-0152
Email: nrspubs@fs.fed.us

Visit our website:
www.nrs.fs.fed.us

Map Disclaimer

United States Department of Agriculture, Forest Service has developed this information for the guidance of its employees, its contractors, and its cooperating Federal, State and other agencies. It is not responsible for the interpretation or use of this information by anyone except its own employees. The use of trade, firm, corporation, or association names in this publication is for the information and convenience of the reader and does not constitute an endorsement by the Department of any product or service. This product is reproduced from geospatial information prepared by the U.S. Department of Agriculture, Forest Service and other U.S. Federal agencies. Geographic Information System (GIS) data and product accuracy may vary. They may be developed from sources of differing accuracy, accurate only at certain scales, based on modeling or interpretation, or incomplete while being created or revised. Using GIS products for purposes other than those for which they were created may yield inaccurate or misleading results. This information was released on the indicated date. The Forest Service reserves the right to correct, update, modify, or replace GIS products without notification.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410, or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.